

# ERYSENG® PARVO/LEPTO

## Suspension for injection for pigs

### STATEMENT OF ACTIVE SUBSTANCES:

One dose (2 ml) contains:

- Inactivated Porcine Parvovirus, strain NADL-2 RP > 1.15\*
- Inactivated *Erysipelothrix rhusiopathiae*, strain R32E11 ELISA > 3.34 IE<sub>50%</sub> \*\*
- Inactivated *Leptospira interrogans* serovar *icterohaemorrhagiae* ≥ 10<sup>8</sup> microorg.
- Inactivated *Leptospira interrogans* serovar *canicola* ≥ 10<sup>8</sup> microorg.
- Inactivated *Leptospira interrogans* serovar *pomona* ≥ 10<sup>8</sup> microorg.
- Inactivated *Leptospira interrogans* serovar *hardjo* ≥ 10<sup>8</sup> microorg.
- Inactivated *Leptospira interrogans* serovar *grippityphosa* ≥ 10<sup>8</sup> microorg.
- Inactivated *Leptospira interrogans* serovar *bratislava* ≥ 10<sup>8</sup> microorg.

\* RP, Relative Potency (ELISA)

\*\* IE<sub>50%</sub> (Inhibition ELISA 50%)

### INDICATIONS:

For the active immunisation of pigs for the protection of progeny against transplacental infection caused by Porcine Parvovirus.

For the active immunisation of pigs to prevent clinical signs and lesions caused by Erysipelas, serotype 1 and serotype 2.

For the active immunisation of pigs to prevent leptospirosis.

### CONTRAINDICATIONS:

Do not use in case of hypersensitivity to the active substances, to the adjuvants or to any of the excipients.

### ADVERSE REACTIONS:

Slight inflammation at the injection site, that lasts for less than 48 hours in most cases.

A very slight and transient increase in body temperature during the first 6 hours after injection, which spontaneously resolves within 24 hours without compromising animal health status.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

### TARGET SPECIES:

Pigs.

### DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:

Intramuscular use.

Administer one dose of 2 ml by intramuscular injection in the neck muscles according to the following schedule:

#### Basic vaccination:

Pigs from 6 months of age which have not been previously vaccinated with the product should be given two injections with an interval of 3-4 weeks. The second injection should be administered 3-4 weeks before mating.

#### Revaccination:

A single injection should be given 2-3 weeks prior to each subsequent mating.

### ADVICE ON CORRECT ADMINISTRATION:

Allow the vaccine to reach room temperature (15-25 °C) before administration.

Shake well before use.

### WITHDRAWAL PERIOD:

Zero days.

### SPECIAL STORAGE PRECAUTIONS:

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C to 8 °C). Do not freeze. Protect from light.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP.

Shelf life after first opening the container: use immediately.

### SPECIAL WARNING(S):

#### Special precautions for use in animals

Vaccinate only healthy animals.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals

In case of adverse reactions following accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

#### Pregnancy and lactation

Can be used during pregnancy and lactation.

#### Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

#### Overdose (symptoms, emergency procedures, antidotes)

No adverse reactions other than already mentioned under section "Adverse reactions" can be expected after the administration of a 2-fold vaccine dose.

#### Incompatibilities

Do not mix with any other veterinary medicinal product.

### SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

### OTHER INFORMATION:

#### Pack sizes:

- Cardboard box with 1 bottle of 10 doses (20 ml).
  - Cardboard box with 1 bottle of 15 doses (30 ml).
  - Cardboard box with 1 bottle of 25 doses (50 ml).
  - Cardboard box with 1 bottle of 50 doses (100 ml).
- Not all pack sizes may be marketed.

Under veterinary prescription

FOR ANIMAL TREATMENT ONLY



Manufactured by HIPRA SAÚDE ANIMAL LTDA. Porto Alegre-RS (Brasil) for:  
LABORATORIOS HIPRA, S.A. Avda. la Selva, 135 . 17170 Amer (Girona) Spain

Tel. (34) 972 43 06 60. Fax (34) 972 43 06 61. hipra@hipra.com